

PHARMACOLOGY AND CELL METABOLISM

No Effect of Prenatal Alcohol Exposure on Activity in Three Inbred Strains of Mice

Chris Downing^{1,*}, Christina Balderrama-Durbin¹, Jonathan Hayes¹, Thomas E. Johnson¹ and David Gilliam²

¹Institute for Behavioral Genetics, University of Colorado, Campus Box 447, Boulder, CO, USA ²Department of Psychological Sciences, University of Northern Colorado, CO, USA

*Corresponding author: 1480 30th St., Room 101, Boulder, CO 80303, USA. Tel: +1-303-492-2152; Fax: +1-303-492-8063; E-mail: cdowning@colorado.edu

(Received 8 February 2008; first review notified 3 April 2008; in revised form 14 July 2008 and 5 August 2008; accepted 2 September 2008; advance access publication 14 October 2008)

Abstract — Aims: Prenatal exposure to alcohol can have adverse effects on the developing fetus. Two of the hallmarks of children exposed to alcohol prenatally are attention deficits and hyperactivity. While hyperactivity has been observed in rats following prenatal ethanol exposure, few studies have examined these effects in mice. The present study investigated the effects of prenatal ethanol exposure on activity in mice from three inbred strains: C57BL/6 (B6), Inbred Long Sleep (ILS) and Inbred Short Sleep (ISS). **Methods:** On Days 7 through 18 of gestation, mice were intragastrically intubated twice daily with either 3.0 g/kg ethanol (E) or an isocaloric amount of maltose–dextrin (MD); non-intubated control (NIC) litters were also generated. Offspring activity was monitored at 30, 60, 90 and 150 days of age. **Results:** While results showed no effects of prenatal ethanol exposure on any measures of activity, we did observe differences in baseline activity among the strains. ISS mice were more active than B6 and ILS for all activity measures except stereotypy; B6 mice had higher measures of stereotypy than ILS and ISS. Younger mice were more active than older mice. The only sex effects were on measures of stereotypy, where males had higher scores. **Conclusions:** Mice are an excellent organism to study genetic influences on many phenotypes. However, our study and others have shown few effects of prenatal ethanol exposure on behavior in mice. It appears as if the prenatal period in mice, corresponding to organogenesis, is not a sensitive period for producing behavioral deficits following ethanol exposure. It is likely that the first 2 weeks postnatally, corresponding to the brain growth spurt, are more sensitive for producing behavioral effects.

INTRODUCTION

Women who consume alcohol (ethanol) during pregnancy place their offspring at risk for a number of teratogenic outcomes. The most severe cases are diagnosed as fetal alcohol syndrome (FAS), a disorder defined by prenatal and/or postnatal growth retardation, a characteristic pattern of craniofacial abnormalities, and central nervous system dysfunction (Jones and Smith, 1973; Jones *et al.*, 1973; Sokol *et al.*, 2003). Because not all offspring exposed to alcohol prenatally display the full spectrum of FAS symptoms (particularly the facial dysmorphism), the term fetal alcohol spectrum disorders (FASD; Koren *et al.*, 2003; Sokol *et al.*, 2003) has been coined to describe varying degrees of ethanol teratogenesis. The estimated incidence of FASD in the United States is around 1% (Sampson *et al.*, 1997; May and Gossage, 2001). Neurodevelopmental and behavioral deficits associated with FAS/FASD include developmental delay, attention deficits, hyperactivity, learning and memory impairments and diminished impulse control (Kelly *et al.*, 1987; Sampson *et al.*, 1997; Coles, 2001; Sokol *et al.*, 2003; Kvigne *et al.*, 2004).

Hyperactivity and attention deficits are hallmarks of children exposed to alcohol prenatally (Mattson and Riley, 1998). Most studies have relied on naturalistic observations and have reported children (from a few hours after birth through high school) exposed to ethanol prenatally as hyperactive, tremulous, fidgety, restless, always on the go, unable to sit still and irritable (Hanson *et al.*, 1976; Streissguth *et al.*, 1978; Shaywitz *et al.*, 1980; Landesman-Dwyer *et al.*, 1981; Nanson and Hiscock, 1990; Steinhausen *et al.*, 1993; Steinhausen and Spohr, 1998; Kvigne *et al.*, 2004). This hyperactivity can occur in the absence of intellectual impairment, at relatively moderate levels of maternal ethanol consumption, and persist throughout childhood. How to best model these behaviors in a rodent

model is unclear. What behaviors in rats and mice best model ‘fidgety’ or ‘restless’ described in the human condition?

Results of prenatal ethanol exposure on activity in rats have been mixed. While many studies have demonstrated increased activity (Martin *et al.*, 1978; Osborne *et al.*, 1979; Bond, 1981; Ulug and Riley, 1983; Meyer and Riley, 1986; Vorhees and Fernandez, 1986), others have not (Vorhees, 1989; Wigal and Amsel, 1990; Westergren *et al.*, 1996; Carneiro *et al.*, 2005). Results depend upon the pattern of ethanol administration, age at which offspring are tested and apparatus in which activity is measured. Most studies used activity monitors (automated or observer scored) and reported total distance traveled, number of photocell beam interruptions or number of squares entered/crossed.

Somewhat surprisingly, few studies have examined hyperactivity in mice exposed to ethanol prenatally. Randall and colleagues investigated the effects of *in utero* ethanol exposure on activity in C57BL/6J (B6) inbred mice. Their paradigms involved giving pregnant dams a liquid diet containing various ethanol concentrations on Days 5/6 through 17/18 of gestation. They reported greater activity in male and female offspring exposed to ethanol prenatally at 23 days of age (Randall *et al.*, 1986), no effect of prenatal ethanol exposure on activity (male and female offspring) at 12, 20 or 36 days (Middaugh *et al.*, 1988) and greater activity in female offspring exposed to ethanol prenatally when tested at 30 days of age (Becker and Randall, 1989). Mothes *et al.* (1996) reported increased activity (36–42 days of age) in the home cage, but not in an activity monitor, in B6 mice exposed to ethanol on days 14–18 of gestation. Allan *et al.* (2003) kept B6SJL/F₁ dams on a liquid ethanol diet before and during pregnancy and reported no effects on offspring activity at 60–100 days. Finally, Gilliam and colleagues (Gilliam *et al.*, 1987; Gilliam, 1990) showed either no effect of prenatal ethanol exposure on activity in Long-Sleep

(LS) or Short-Sleep (SS) mice (21–25 days of age), or an increase in activity in SS mice and mice derived from reciprocal crosses between SS and LS (150 days of age) exposed to ethanol prenatally.

Studies have shown that, in humans, genetic factors can influence susceptibility and resistance to ethanol teratogenesis. Several case study reports indicate that monozygotic twins are more similarly affected than dizygotic twins (Palmer *et al.*, 1974; Christoffel and Salafsky, 1975; Chasnoff, 1985; Riikonen, 1994). A more comprehensive study examined ethanol exposure *in utero* in both monozygotic and dizygotic twins. The rate of concordance for diagnosis was 5/5 for monozygotic twins and 7/11 for dizygotic twins, and the authors concluded that genes had a modulating influence on expression of the teratogenic effects of alcohol (Streissguth and Dehaene, 1993). More recently, several studies have shown that different alleles of the alcohol dehydrogenase gene (ADH), an enzyme involved in ethanol metabolism, can influence the severity of teratogenesis in different ethnic populations (McCarver *et al.*, 1997; Viljoen *et al.*, 2001; Stoler *et al.*, 2002; Das *et al.*, 2004). Therefore, characterizing mice for hyperactivity following prenatal ethanol exposure (or any prenatal alcohol trait) is important because mice are a much more tractable species than rat in which to examine genetics. Many more inbred, recombinant inbred, congenic and selectively bred lines of mice exist than rat. In addition, the mouse genome is much better annotated and many more strains of mice have been sequenced. Finally, it is currently much easier to manipulate the mouse genome (i.e. targeted mutagenesis) than the rat genome.

In the present study, we examined activity following *in utero* ethanol exposure in mice from three inbred strains: Inbred Long-Sleep (ILS), Inbred Short-Sleep (ISS) and B6. We chose these strains because they are among the few mouse strains that have been characterized for any prenatal alcohol phenotypes. The ILS and ISS mice were derived from LS and SS mice, selectively bred for differential sensitivity to a hypnotic dose of alcohol (McClearn and Kakihana, 1981). Previous research has shown that LS mice are more susceptible than SS mice to several measures of teratogenesis, including activity as noted above (Gilliam *et al.*, 1989a, 1989b; Gilliam, 1990; Gilliam and Kotch, 1990, 1996). This suggests that one or more genes that mediate differential sensitivity to the hypnotic effects of ethanol in SS and LS may also mediate differential sensitivity to ethanol teratogenesis. The ILS and ISS mice have not been characterized for any prenatal alcohol phenotypes.

B6 mice are relatively susceptible to fetal weight deficits and kidney, limb and skeletal malformations following prenatal ethanol exposure (Webster *et al.*, 1980; Gilliam and Irtenkauf, 1990; Boehm *et al.*, 1997; Gilliam *et al.*, 1997; Downing and Gilliam, 1999). As discussed above, results are mixed when B6 mice are examined for hyperactivity following prenatal ethanol exposure.

METHODS

Animals

Male and female ILS, ISS and B6 mice were obtained from and housed in the specific pathogen-free (SPF) facility at the Institute for Behavioral Genetics, Boulder, CO, USA. Males were individually housed while females were housed 3–5 per

cage; mice were maintained on a 12-h light/dark cycle (lights on at 7:00 am) and were given food and water *ad libitum*. The temperature was kept at a constant 22°C. All procedures were approved by the University of Colorado Institutional Animal Care and Use Committee, in accordance with National Institute of Health guidelines.

Mating and dosing

Females weighed a minimum of 18 g at mating. Two females were placed in each male's cage overnight and examined for a seminal plug in the morning as evidence of mating. Plugged females were weighed, randomly assigned to a treatment condition (ethanol, E; maltose–dextrin, MD; or non-intubated control, NIC) and single-housed. The day of plug detection was designated as Day 0 of pregnancy (gestational day 0: GD 0). On GD 7, females were weighed to ascertain a 1.5 g minimum weight gain. Pregnant dams were then intubated twice daily (9:00 am and 3:00 pm, GD 7–GD 18) with either 3.0 g/kg ethanol (20% w:v) or an isocaloric amount of MD. Mice in the NIC group were weighed daily; this group served as a control to assess the effects of repeated handling and intubations in the E and MD groups.

Dams were checked for births twice daily starting on GD 19. Once born (postnatal day 0; PND 0), litters were weighed and sexed. Litters were weighed again on PND 3 and offspring sex confirmed. Litters were then culled to four offspring, two males and two females when possible. Offspring were again weighed on PNDs 5, 10, 20 and subsequent days when activity testing took place. Pups were weaned on PND 28 and housed with same-sex mice, 4–5 per cage. Mothers who died during the intubation process or failed to deliver by GD 20 were sacrificed and the number of implantation sites was counted.

Offspring testing

Offspring were tested for activity in 2-day intervals on PNDs 30 and 31, 60 and 61, 90 and 91, and 150 and 151. We chose the 30- and 60-day timepoints because they represent a reasonable approximation of the adolescent and young adult stages in mice. We also tested offspring at 90 and 150 days to see if effects persisted into adulthood; Gilliam (1990) found effects of prenatal ethanol exposure on activity in offspring tested at 150 days of age. Testing took place between 9:00 am and 11:00 am. Mice were placed in an automated activity monitor for 15 min (three successive 5-min bins). The activity monitoring system (Digiscan: Accuscan Instruments Inc., Columbus, OH, USA) consists of a 16-beam photocell apparatus interfaced with a PC; the monitors consist of a 40-cm² chamber with a 30.5 cm ceiling. Software interfacing with the monitors records the total photocell beam breaks, both vertical and horizontal. Computer software recorded a number of other activity measures, as described below.

Maternal blood ethanol concentration

A separate group of pregnant females were examined for blood ethanol concentration (BEC). Dams were intubated twice daily with 3.0 g/kg ethanol from GD 7 to GD 18, as described above. Blood was drawn from the retro-orbital sinus 30, 60, 120 and 180 min after the second intubation (3:00 pm) on GD 18. Ten microliters of retro-orbital blood was added to 200 μ l of perchloric acid on ice to precipitate blood solids. Blood samples were vortexed and centrifuged at 4500 rpm for 10 min.

Table 1. Mean (\pm SEM) percent maternal weight gain, litter size, prenatal mortality and pup weight at birth

	C57BL/6			ILS			ISS		
	E (15)	MD (9)	NIC (12)	E (7)	MD (7)	NIC (8)	E (11)	MD (9)	NIC (9)
% Wt gain ^a	45 ^b (4)	67 (3)	69 (5)	49 (3)	44 (5)	65 ^c (2)	30 (2)	30 (5)	33 (4)
Lit size ^d	6.20 (0.75)	7.78 (0.72)	7.60 (0.43)	5.29 (0.42)	5.57 (0.90)	6.88 (0.40)	3.18 (0.40)	3.38 (0.32)	3.67 (0.37)
% PM ^e	56 (12)	22 (12)	14 (7)	4 (4)	32 (18)	8 (4)	22 (12)	15 (7)	11 (11)
Pup Wt. ♀	1.23 (0.05)	1.37 (0.04)	1.34 (0.04)	1.34 (0.05)	1.49 (0.06)	1.36 (0.03)	1.53 (0.03)	1.60 (0.05)	1.59 (0.04)
Pup Wt. ♂	1.33 (0.05)	1.33 (0.03)	1.33 (0.06)	1.51 (0.08)	1.47 (0.05)	1.46 (0.03)	1.53 (0.04)	1.86 (0.13)	1.67 (0.06)

^aPercent maternal weight gain calculated as: weight on GD 18 – weight on GD 7/weight on GD 7.

^bEthanol-treated B6 dams put on less weight than MD and NIC controls, $P_s < 0.01$.

^cEthanol- and maltose-treated ILS dams put on less weight than NIC controls, $P_s < 0.03$ and 0.01 , respectively.

^dB6 and ILS litters were larger than ISS litters, $P_s < 0.001$.

^ePostnatal mortality calculated as: No. of pups on PND 0 – No. of pups on PND 3 before culling/No. of pups on PND 0.

The plasma or supernatant was then removed from the pellet and an equal volume of KOH was added to the supernatant to neutralize the perchloric acid. The sample was then vortexed and stored in the freezer until analysis (once per week). BEC was determined by spectrophotometric analysis of an enzyme assay as described by Smolen *et al.* (1986).

Statistical analyses

Data were examined using analysis of variance (ANOVA) with strain (ILS, ISS and B6), treatment (E, MD, NIC) and sex as grouping factors. For maternal data, percent weight gain during pregnancy, litter size and postnatal mortality were examined. For offspring data, in order to control for litter effects and inflated sample size, litter means were the unit of analyses (Zorilla, 1997; Wainwright, 1998). Offspring weight at birth was examined using ANOVA. Body weight and activity measures from PNDs 3–151 were examined with repeated measures ANOVA (RM-ANOVA), with strain, treatment and sex as between-group factors and age as the within-group factor. When significant main effects or interactions were observed, *post hoc* analyses were done using Bonferroni corrected *t*-tests.

The following activity variables were analyzed: horizontal activity, total distance traveled (cm), horizontal movement number, movement time, rest time, vertical movement number, vertical time, stereotypy number and stereotypy time. As noted earlier, most previous studies have simply looked at total distance traveled and number of beam breaks. Examining these additional activity variables should provide insights into what measures in mice may best model ‘hyperactivity’ in children exposed to ethanol *in utero*.

Activity data was analyzed in several ways. First, we analyzed data within each session. For almost every day and measure, we saw habituation; animals were less active in the last 5-min bin compared to the first 5-min bin. There were no effects of genotype, treatment or sex on habituation. We also analyzed 15-min totals and found very few differences compared to individual 5-min bins; therefore, we present data from 15-min totals only. Data from the first day of each 2-day session was analyzed separately (30, 60, 90 and 150), from the second day of each 2-day session (31, 61, 91, 151) and all days (30, 31, 60, 61, 90, 91, 150, 151). While activity measures were generally lower when the second day was analyzed separately (likely habituation), the pattern of results was remarkably consistent. Thus, we present activity analyses (RM-ANOVA) from all days. Figures include means for 2-day averages (30 and 31, 60 and 61, etc.).

RESULTS

Maternal data

Resorptions. One out of 11 ISS MD and 1 out of 14 ISS E dams lost their litters (did not give birth and had resorptions when sacrificed on GD 20). While no ILS dams lost their litters, 3 of 11 E dams and 1 of 9 MD dams died following intubation. No B6 females died during intubations or lost their litters.

Weight gain, litter size and postnatal mortality. ANOVA showed significant main effects of strain ($F(2, 78) = 42.45$, $P < 0.001$) and treatment ($F(2, 78) = 8.36$, $P < 0.01$) and a significant strain by treatment interaction ($F(4, 78) = 4.16$, $P < 0.01$) for percent maternal weight gain (Table 1). *Post hoc* analyses showed that ISS dams put on less weight than ILS and B6 ($P_s < 0.01$). As expected, NIC dams gained more weight than E and MD treatment groups ($P_s < 0.01$). Decomposition of the strain by treatment interaction showed that E-treated B6 females put on significantly less weight than MD and NIC controls ($P_s < 0.01$). In addition, E- and MD-treated ILS females put on less weight than their NIC controls ($P_s < 0.03$ and 0.01 , respectively).

While there was a main effect of strain ($F(2, 80) = 32.83$, $P < 0.001$), there was no effect of treatment and no strain by treatment interaction on litter size. *Post hoc* analysis showed that B6 and ILS dams had larger litters than ISS dams ($P_s < 0.001$; Table 1). It should be noted that litter size is likely an underestimate. On GD 19, we examined dams for litters twice, in the morning and afternoon. In several cases, when litters were found, the dams had partially eaten one or more pups; these pups were included in our ‘litter size’ variable and accounted for some of the postnatal mortality. It seems likely that there were a few dams that had completely eaten pups by the time litters were found, so litter size is likely underestimated.

Postnatal mortality was calculated as: (number of pups on PND 0 – number of pups on PND 3 before culling)/number of pups on PND 0. There were no main effects of strain or treatment on postnatal mortality.

Blood ethanol concentration. Five pregnant dams per genotype were intubated with 3.0 g/kg ethanol, twice daily, from GD 7 to GD 18. Thirty, 60, 120 and 180 min following the last intubation on GD 18 (3:00 pm), blood was obtained and assayed for ethanol concentration. Blood ethanol levels averaged 317 mg/dl 30 min after the last injection and declined to 56–117 mg/dl at 180 min (Fig. 1). Data were analyzed using RM-ANOVA, with time after intubation (30, 60, 120, 180 min) as the within-group variable and strain as the between-group




Fig. 1. Mean (SEM) blood ethanol concentration for pregnant C57BL/6J (B6), Inbred Long-Sleep (ILS) and Inbred Short-Sleep (ISS) mice. Blood was drawn 30, 60, 120 and 180 minutes following the last intubation on GD 18. RM-ANOVA showed only a main effect of timepoint ($P < 0.01$).

variable. Results showed a significant main effect of time ($F(3, 30) = 288.65$, $P < 0.01$) but no significant main effect of strain and no significant strain by time interaction. In all strains, BECs declined in a similar manner across time.

Offspring data

Birth weight. Birth weight and offspring weight from PNDs 3–151 were analyzed using litter means for each sex as the unit of analysis. We found significant main effects of strain ($F(2, 119) = 47.49$, $P < 0.001$), treatment ($F(2, 119) = 4.87$, $P < 0.01$) and sex ($F(1, 119) = 6.82$, $P < 0.02$) for offspring weight at birth but no significant interactions among the variables. *Post hoc* analyses showed that ISS offspring weighed more than ILS and B6 (P s < 0.001) and ILS offspring weighed more than B6 ($P < 0.01$); MD-treated litters weighed more than E-treated litters ($P < 0.01$) and NIC litters ($P < 0.02$); males weighed more than females ($P < 0.02$; Table 1).

Weight gain. Offspring weight from PNDs 3–151 was analyzed using RM-ANOVA, with age as a within-subjects variable and strain, treatment and sex as between-subjects variables. ANOVAs involving repeated measures used the Greenhouse–Geisser adjustment factor to assess the significance of the observed F ratio. We found significant main effects of age ($F(3.79, 378) = 1268$, $P < 0.001$), strain ($F(2, 100) = 23.32$, $P < 0.001$) and sex ($F(1, 100) = 189.05$, $P < 0.001$) on weight gain from PNDs 3–151 but no main effect of treatment. Age interacted with strain ($F(7.57, 378) = 23.67$, $P < 0.001$) and sex ($F(3.78, 378) = 186.08$, $P < 0.001$) and there was a significant age by strain by sex interaction ($F(7.57, 378) = 2.43$, $P < 0.02$). Simple effects analysis (within strain) at each age showed that in general, ISS offspring gained more weight than the other three strains from PNDs 3–20, while ILS offspring gained less weight than the other three strains from PNDs 3–60; males gained more weight than females from PNDs 20–151 (P s < 0.001).

Measures of activity. Total distance traveled (TDT) and horizontal activity (HACT) are the two most commonly used measures of behavioral activation in rodents. TDT indicates, in




Fig. 2. Mean (SEM) total distance traveled (TDT) for B6, ILS and ISS mice. Bars represent the average of 2 days of testing. For example, Day 30 equals the average TDT on Days 30 and 31. For Figs. 2–5, sample sizes (litters): B6 E = 13, B6 MD = 9, B6 NIC = 11; ILS E = 7, ILS MD = 7, ILS NIC = 8; ISS E = 10, ISS MD = 9, ISS NIC = 9. *ISS mice were more active than ILS and B6 on Days 30, 60 and 90; P s < 0.05 . **ILS were more active than B6 at 150 days; $P < 0.01$.

centimeters, the distance traveled during a given sample period, while HACT is the total number of beam breaks that occur in the horizontal plane during a given sample period. For TDT, RM-ANOVA showed significant main effects of age ($F(3.84, 421) = 12.55$, $P < 0.001$) and strain ($F(2, 110) = 34.54$, $P < 0.001$). There was no effect of treatment. *Post hoc* analyses showed that ISS offspring had greater TDT (Fig. 2) compared to ILS and B6 (P s < 0.01); younger mice had a higher TDT than older mice (P s < 0.01). RM-ANOVA also showed a significant age by strain interaction for TDT ($F(7.67, 421) = 7.45$, $P < 0.001$). ISS mice had higher TDT (Fig. 2) than B6 and ILS at 30, 60 and 90 days of age (P s < 0.05). We found identical results (with slightly different F -values; data not shown) for HACT. In general, all strains had a decrease in activity across days. This could reflect habituation or perhaps mice simply are not as active when they get older. Interestingly, the exception to this characterization is ILS mice, which had a large increase in activity at 150 days.

Movement number (MOVNO) is the number of separate horizontal movements an animal makes in a given sample period. Movement time (MOVTIM) is the amount of time an animal ambulates (horizontally) in a given sample period while rest time (RSTIM) is the amount of time an animal does not ambulate. For all three variables, RM-ANOVA showed significant main effects of age and strain, and a significant age by strain interaction (all P s < 0.001). There was no effect of treatment. As can be seen in Fig. 3, ISS made more movements and spent more time ambulating than B6 and ILS (P s < 0.01). Because RSTIM is simply the inverse of MOVTIM, we present data from MOVTIM only.

We examined two measures of vertical activity. Vertical movement number (VMVNO) is the number of movements an animal makes in the vertical plane (rearing), while vertical time (VTIM) is the amount of time an animal spends rearing. For VMVNO, ANOVA showed significant main effects of age ($F(5.47, 601) = 22.38$, $P < 0.01$) and strain ($F(2, 110) = 24.09$,




Fig. 3. Mean (SEM) movement number (MVNO) and movement time (MVTIM) in B6, ILS and ISS mice. Each timepoint represents a 2-day average. *ISS mice had greater MVNO than ILS at 30, 60 and 90 days of age ($P_s < 0.001$); ISS had greater MVNO than B6 at 30 and 90 days ($P_s < 0.001$). **ISS also had greater MVTIM than ILS at 30, 60 and 90 days ($P_s < 0.001$); they also had greater MVTIM than B6 on all days tested ($P_s < 0.001$).

$P < 0.01$) and a significant age by strain interaction ($F(10.94, 601) = 13.22$, $P < 0.01$). *Post hoc* analyses showed that ILS offspring made fewer vertical movements than B6 and ISS ($P_s < 0.01$; Fig. 4). All three strains had a significant increase in vertical movements at the 90-day timepoint compared to the other timepoints ($P_s < 0.01$; Fig. 4). For VTIM, ANOVA again showed significant main effects of age ($F(5.14, 565) = 69.15$, $P < 0.01$) and strain ($F(2, 110) = 42.98$, $P < 0.01$) and a significant age by strain interaction ($F(10.28, 565) = 16.46$, $P < 0.01$). *Post hoc* analyses showed that ISS offspring had significantly greater VTIM compared to B6 and ILS ($P_s < 0.01$; Fig. 4). There were no treatment effects on either variable.

If a mouse repeatedly breaks the same beam or set of beams, the mouse is exhibiting stereotypic behavior. We analyzed two measures of stereotypic behavior. Stereotypy number (STNO) is the number of times the monitor observed stereotypic behavior in the mouse; a break in stereotypy of one second or more is required to separate one stereotypic episode from the next. Stereotypy time (STIM) is the total amount of time that stereotypic behavior is exhibited. For STNO, RM-ANOVA showed significant main effects of age ($F(5.27, 579) = 25.49$, $P < 0.01$), strain ($F(2, 110) = 40.28$, $P < 0.01$) and sex ($F(1, 100) = 6.91$, $P < 0.01$) and significant age by strain ($F(10.53, 579) = 6.30$, $P < 0.01$) and strain by sex ($F(2, 110) = 3.97$, $P < 0.03$) interactions. *Post hoc* analyses showed that B6 offspring had




Fig. 4. Mean (SEM) vertical movement number and vertical movement time in B6, ILS and ISS mice. Each timepoint represents a 2-day average. *ILS had less VMOV than ISS and B6 at 30 and 60 days ($P < 0.01$); **ILS had less VTIM than ISS and B6 at 30 and 60 ($P < 0.01$); ***ISS had greater VTIM than B6 at 60 ($P < 0.05$), 90 and 150 days ($P < 0.01$).

significantly greater STNO than ILS and ISS, while ILS offspring had significantly less STNO than B6 and ISS (all $P < 0.01$). In addition, males had significantly more STNO than females at 60, 90 and 150 days of age ($P < 0.05$). As can be seen in Fig. 5, the effect of sex was strain and age dependent. Similarly, for STIM, ANOVA showed significant main effects of age ($F(5.01, 551) = 34.63, P < 0.01$), strain ($F(2, 110) = 46.08, P < 0.01$) and sex ($F(1, 110) = 5.97, P < 0.02$). Age also interacted with strain ($F(10.02, 551) = 10.25, P < 0.01$) and sex ($F(5.01, 551) = 2.32, P < 0.05$). *Post hoc* analyses showed that, similar to STNO, B6 had significantly greater STIM than

ILS and ISS while ILS had significantly less STIM than B6 and ISS; males had significantly higher STIM than females (all $P < 0.01$). There was no effect of treatment on STNO or STIM.

DISCUSSION

Attention deficits and hyperactivity are hallmarks of children exposed to ethanol *in utero*. While the effects of prenatal ethanol exposure on attention and activity have been fairly well characterized in rats, few studies have examined these effects in mice.




Fig. 5. Mean (SEM) stereotypy number (STNO) in B6 (a), ILS (b) and ISS (c) mice. Each timepoint represents a 2-day average. *Males had significantly greater STNO than females, $P < 0.01$. **Males had significantly greater STNO than females, $P < 0.05$.

For activity, while two studies have reported increased activity in B6 mice exposed to ethanol prenatally (Randall *et al.*, 1986; Becker and Randall, 1989), two others have not (Middaugh *et al.*, 1988; Mothes *et al.*, 1996). Allan *et al.* (2004) reported no effects of prenatal ethanol exposure on activity in B6 \times SJL F₁ mice. Results from our study showed no effects of prenatal ethanol exposure on any measures of activity in B6 offspring.

LS and SS mice were selectively bred for sensitivity (LS) and resistance (SS) to the soporific effects of ethanol, as measured by loss of the righting reflex (LORE: loss of righting due to ethanol; McClearn and Kakihana, 1981). ILS and ISS were derived by subsequent inbreeding of LS and SS. In addition to the hypnotic effects of ethanol, LS, SS, ILS and ISS differ on many other behavioral and physiological traits. Compared to LS and ILS, SS and ISS mice have higher baseline activity and also show greater activation following a low dose of ethanol (Dudek and Abbott 1984; Phillips and Dudek, 1991; Owens *et al.*, 2002). In addition, these lines of mice differ in measures of functional tolerance following ethanol exposure (Gill and Deitrich, 1998; Deitrich *et al.*, 2000; Bennett *et al.*, 2007) and in their responses to other sedative hypnotics (Christensen *et al.*, 1996; Simpson *et al.*, 1998). Purkinje cells of the cerebellum of LS mice are also much more sensitive to the depressant effects of ethanol than Purkinje cells of SS mice, as determined by electrophysiological measures (Sorensen *et al.*, 1980; Basile *et al.*, 1983). Levels of aldehyde dehydrogenase in Purkinje cells of the cerebellum are higher in SS mice compared to LS mice (Zimatkin and Deitrich, 1995). Purkinje cells of the cerebellum are particularly vulnerable to neonatal (third trimester equivalent) ethanol exposure in rats (Goodlett *et al.*, 1998; Thomas *et al.*, 1998a; Light *et al.*, 2002).

LORE is a measure of initial sensitivity to alcohol. Previous research suggested that individual sensitivity to alcohol might influence susceptibility and resistance to some of the detrimental effects of prenatal alcohol exposure. Ethanol-sensitive LS mice are susceptible to fetal weight deficits, postnatal growth deficits, increased postnatal mortality, fetal brain weight reductions and skeletal malformations following prenatal alcohol exposure, while ethanol-insensitive SS mice are relatively resistant (Gilliam *et al.*, 1989a, 1989b; Goodlett *et al.*, 1989; Gilliam and Irtenkauf, 1990; Gilliam and Kotch, 1990, 1996). In addition, LS mice exposed to ethanol prenatally took significantly more trials to reach a passive avoidance criterion than their controls, while SS mice did not. Two studies looked at hyperactivity in LS and SS mice following *in utero* ethanol exposure. Gilliam *et al.* (1987) reported no effects of prenatal ethanol exposure on activity in young LS and SS mice. In contrast, Gilliam (1990) reported an increase in activity at 150 days of age in SS, but not LS, mice exposed to alcohol prenatally. This increase in activity at 150 days was not confirmed in ISS mice in the present study. Two major differences between the present study and the two Gilliam studies are that mice were cross-fostered at birth and were tested in sound attenuation chambers under red light.

ISS showed greater baseline activity than ILS on all measures. The differences in TDT and HACT verify results from previous studies (see above) and show that these differences exist at younger ages than previously reported. This suggests that in addition to LORE, differences in baseline activity were also selected for in LS and SS, and were captured during the inbreeding process. B6 were intermediate on most measures of activity except for our two measures of stereotypy, where B6 were higher than both ISS and ILS. The only sex differences in baseline activity were also for stereotypy, where males displayed higher rates than females.

Few studies have demonstrated effects on behavior in mice following prenatal alcohol exposure. This is likely due, at least in part, to behavior being correlated with brain

development/functioning. This is one area where humans and mice differ. In mammals, a period of rapid central nervous system growth and proliferation (the 'brain growth spurt') occurs during the third trimester. While the third trimester of pregnancy occurs *in utero* in humans, it occurs during approximately the first 2 weeks postnatally in rodents (Dobbing and Sands, 1979). Thus, in order to mimic the third trimester ethanol exposure in rodents, one must administer ethanol to neonatal pups. Only a handful of studies have examined behavior (including hyperactivity) in mice following early postnatal ethanol exposure (Yanai and Ginsburg, 1977, 1979; Yanai, 1983; Ciociola and Gautieri, 1988; Pick *et al.*, 1993; Pal and Alkana, 1997; Wozniak *et al.*, 2004). When exposed to ethanol neonatally, rats reliably show changes in behavior (Goodlett *et al.*, 1987; Kelly *et al.*, 1987; Melcer *et al.*, 1994; Pauli *et al.*, 1995; Goodlett and Johnson, 1997; Thomas *et al.*, 1998b, 2003). Therefore, researchers using mice as a model for the behavioral effects of FASD should consider early postnatal exposure.

Acknowledgements — This study was supported by grants AA016676 and AA01466.

REFERENCES

- Allan AM, Chynoweth J, Tyler LA *et al.* (2003) A mouse model of prenatal ethanol exposure using a voluntary drinking paradigm. *Alcohol Clin Exp Res* **27**:2009–16.
- Basile A, Hoffer B, Dunwiddie T. (1983) Differential sensitivity of cerebellar Purkinje neurons to ethanol in selectively bred outbred lines of mice: maintenance *in vitro* independent of synaptic transmission. *Brain Res* **264**:69–78.
- Becker HC, Randall CL (1989) Effects of prenatal ethanol exposure in C57BL mice on locomotor activity and passive avoidance. *Psychopharmacology* **97**:40–4.
- Bennett B, Downing C, Carosone-Link P *et al.* (2007) Quantitative trait locus mapping for acute functional tolerance to ethanol in the L \times S recombinant inbred panel. *Alcohol Clin Exp Res* **31**:1–9.
- Boehm SL, Lundahl KR, Caldwell J *et al.* (1997) Ethanol teratogenesis in the C57BL/6J, DBA/2J and A/J inbred mouse strains. *Alcohol* **14**:389–95.
- Bond NW. (1981) Prenatal alcohol exposure in rodents: a review of its effects on offspring activity and learning ability. *Aust J Psych* **33**:331–44.
- Carneiro LMV, Diogenes JPL, Vasconcelos SMM *et al.* (2005) Behavioral and neurochemical effects on rat offspring after prenatal exposure to ethanol. *Neurotoxicol Teratol* **27**:585–92.
- Chasnoff IJ. (1985) Fetal alcohol syndrome in twin pregnancy. *Acta Genet Med Gemellol (Roma)* **34**:229–32.
- Christensen SC, Johnson TE, Markel PD *et al.* (1996) Quantitative trait locus analyses of sleep-times induced by sedative-hypnotics in LS \times SS recombinant inbred strains of mice. *Alcohol Clin Exp Res* **20**:543–50.
- Christoffel KK, Salafsky I. (1975) Fetal alcohol syndrome in dizygotic twins. *J Pediatr* **87**:963–7.
- Ciociola AA, Gautieri RF. (1988) Teratogenic and behavioral anomalies induced by acute exposure of mice to ethanol and their possible relation to fetal brain DNA synthesis. *Pharm Res* **5**:447–52.
- Coles CD. (2001) Fetal alcohol exposure and attention: moving beyond ADHD. *Alcohol Res Health* **25**:199–203.
- Das UG, Cronk CE, Martier SS *et al.* (2004) Alcohol dehydrogenase 2 \times 3 affects alterations in offspring facial dysmorphology associated with maternal ethanol intake in pregnancy. *Alcohol Clin Exp Res* **28**:1598–606.
- Deitrich RA, Bludeau P, Erwin VG. (2000) Phenotypic and genotypic relationships between ethanol tolerance and sensitivity in mice selectively bred for initial sensitivity to ethanol (SS and LS) or development of acute tolerance (HAFT and LAFT). *Alcohol Clin Exp Res* **24**:595–604.
- Dobbing J, Sands J. (1979) Comparative aspects of the brain growth spurt. *Early Hum Dev* **3**:79–83.
- Downing C, Gilliam D. (1999) Cytoplasmic factors do not contribute to a maternal effect on ethanol teratogenesis. *Behav Genet* **29**:31–9.
- Dudek BC, Abbott ME. (1984) The relationship between ethanol-induced locomotor activation and narcosis in long-sleep and short-sleep mice. *Alcohol Clin Exp Res* **8**:272–6.
- Gill K, Deitrich RA. (1998) Acute tolerance to the ataxic effects of ethanol in short-sleep (SS) and long-sleep (LS) mice. *Psychopharmacology* **136**:91–8.
- Gilliam DM. (1990) Maternal genetic effects on alcohol behavioral teratology. *Alcohol Clin Exp Res* **14**:293.
- Gilliam DM, Irtenkauf KT. (1990) Maternal genetic effects on ethanol teratogenesis and dominance of relative embryonic resistance to malformations. *Alcohol Clin Exp Res* **14**:539–45.
- Gilliam DM, Kotch LE. (1990) Alcohol-related birth defects in long- and short-sleep mice: postnatal litter mortality. *Alcohol* **7**:483–7.
- Gilliam DM, Kotch LE. (1996) Dose-related growth deficits in LS but not SS mice prenatally exposed to alcohol. *Alcohol* **13**:47–51.
- Gilliam DM, Kotch LE, Dudek BC *et al.* (1989a) Ethanol teratogenesis in mice selected for differences in alcohol sensitivity. *Alcohol* **5**:513–9.
- Gilliam DM, Kotch LE, Dudek BC *et al.* (1989b) Ethanol teratogenesis in selectively bred long-sleep and short-sleep mice: a comparison to inbred C57BL/6J mice. *Alcohol Clin Exp Res* **13**:667–72.
- Gilliam DM, Mantle MA, Barkhausen DA *et al.* (1997) Effects of acute prenatal ethanol administration in a reciprocal cross of C57BL/6J and short-sleep mice: maternal effects and nonmaternal factors. *Alcohol Clin Exp Res* **21**:28–34.
- Gilliam DM, Stilman A, Dudek BC *et al.* (1987) Fetal alcohol effects in long- and short-sleep mice: activity, passive avoidance and *in utero* ethanol levels. *Neurotoxicol Teratol* **9**:349–57.
- Goodlett CR, Gilliam DM, Nichols JM *et al.* (1989) Genetic influences on brain growth restriction induced by developmental exposure to alcohol. *Neurotoxicology* **10**:321–34.
- Goodlett CR, Johnson TB. (1997) Neonatal binge ethanol exposure using intubation: timing and dose effects on place learning. *Neurotoxicol Teratol* **6**:435–46.
- Goodlett CR, Kelly SJ, West JR. (1987) Early postnatal alcohol exposure that produces high blood alcohol levels impairs development of spatial navigation learning. *Psychobiology* **15**:64–74.
- Goodlett CR, Pearlman AD, Lundahl KR. (1998) Binge neonatal alcohol intubations induce dose-dependent loss of Purkinje cells. *Neurotoxicol Teratol* **20**:285–92.
- Hanson JW, Jones KL, Smith DW. (1976) Fetal alcohol syndrome. Experience with 41 patients. *JAMA* **235**:1458–60.
- Jones KL, Smith DW. (1973) Recognition of the fetal alcohol syndrome in early infancy. *Lancet* **2**:999–1001.
- Jones KL, Smith DW, Ulleland CN *et al.* (1973) Pattern of malformations in offspring of chronic alcoholic mothers. *Lancet* **1**:1267–71.
- Kelly SJ, Pierce DR, West JR. (1987) Microencephaly and hyperactivity in adult rats can be induced by neonatal exposure to high blood alcohol concentrations. *Exp Neurol* **96**:580–93.
- Koren G, Nulman I, Chudley AE *et al.* (2003) Fetal alcohol spectrum disorder. *CMAJ* **169**:1181–5.
- Kvigne VL, Leonardson GR, Neff-Smith M *et al.* (2004) Characteristics of children who have full or incomplete fetal alcohol syndrome. *J Pediatr* **145**:635–40.
- Landesman-Dwyer S, Ragozin AS, Little RE. (1981) Behavioral correlates of prenatal alcohol exposure: a four-year follow-up study. *Neurobehav Toxicol Teratol* **3**:187–93.
- Light KE, Belcher SM, Pierce DR. (2002) Time course and manner of Purkinje neuron death following a single ethanol on postnatal day 4 in the developing rat. *Neuroscience* **114**:327–37.
- Martin JC, Martin DC, Sigman G *et al.* (1978) Maternal ethanol consumption and hyperactivity in cross-fostered offspring. *Physiol Psychol* **6**:362–5.
- Mattson SN, Riley EP. (1998) A review of the neurobehavioral deficits in children with fetal alcohol syndrome or prenatal exposure to alcohol. *Alcohol Clin Exp Res* **22**:279–94.
- May PA, Gossage JP (2001) Estimating the prevalence of fetal alcohol syndrome. *Alcohol Res Health* **25**:159–67.

- McCarver DG, Thomasson HR, Martier SS *et al.* (1997) Alcohol dehydrogenase-2*3 allele protects against alcohol-related birth defects among African Americans. *J Pharmacol Exp Ther* **283**:1095–101.
- McClern GE, Kakihana R. (1981) Selective breeding for ethanol sensitivity: short-sleep and long-sleep mice. In McClern GE, Deitrich RA, Erwin VG (eds). *Development of Animal Models as Pharmacogenetic Tools*. Washington, DC: NIAAA Research Monograph 6, US Department of Health and Human Services, 147–59.
- Melcer T, Gonzalez D, Riley EP. (1994) Locomotor activity and alcohol preference in alcohol-preferring and—nonpreferring rats following neonatal alcohol exposure. *Neurotoxicol Teratol* **17**: 41–8.
- Meyer LS, Riley EP. (1986) Behavioral teratology of alcohol. In Riley EP, Vorhees CV (eds). *Handbook of Behavioral Teratology*. New York: Plenum, 101–40.
- Middaugh LD, Randall CL, Favara JP. (1988) Prenatal ethanol exposure in C57 mice: effects on pregnancy and offspring development. *Neurotoxicol Teratol* **10**:175–80.
- Moths HK, Opitz B, Werner R *et al.* (1996) Effects of prenatal ethanol exposure and early experience on home-cage and open-field activity in mice. *Neurotoxicol Teratol* **18**:59–65.
- Nanson JL, Hiscock M. (1990) Attention deficits in children exposed to alcohol prenatally. *Alcohol Clin Exp Res* **14**:656–61.
- Osborne GL, Caul WF, Fernandez (1979) Behavioral effects of prenatal ethanol exposure and differential early experience in rats. *Pharmacol Biochem Behav* **12**:393–401.
- Owens JC, Stallings MC, Johnson TE (2002) Genetic analysis of low-dose ethanol-induced activation (LDA) in inbred long-sleep (ILS) and inbred short-sleep (ISS) mice. *Behav Genet* **32**:163–71.
- Pal N, Alkana RL (1997) Use of inhalation to study the effect of ethanol and ethanol dependence on neonatal mouse development without maternal separation: a preliminary study. *Life Sci* **61**:1269–81.
- Palmer RH, Ouellette EM, Warner L *et al.* (1974) Congenital malformations offspring of a chronic alcoholic mother. *Pediatrics* **53**: 490–4.
- Pauli J, Wilce P, Bedi KS. (1995) Spatial learning ability of rats following acute exposure to alcohol during early postnatal life. *Physiol Behav* **58**:1013–20.
- Phillips TJ, Dudek BC. (1991) Locomotor activity responses to ethanol in selectively bred long- and short-sleep mice, two inbred mouse strains, and their F₁ hybrids. *Alcohol Clin Exp Res* **16**:255–61.
- Pick CG, Cooperman M, Trombka D *et al.* (1993) Hippocampal cholinergic alterations and related behavioral deficits after early exposure to ethanol. *Int J Dev Neurosci* **11**:379–85.
- Randall CL, Becker HC, Middaugh LD. (1986) Effect of prenatal ethanol exposure on activity and shuttle avoidance behavior in adult C57 mice. *Alcohol Drug Res* **6**:351–60.
- Riikonen RS. (1994) Difference in susceptibility to teratogenic effects of alcohol in discordant twins exposed to alcohol during the second half of gestation. *Pediatr Neurol* **11**:332–6.
- Sampson PD, Streissguth AP, Bookstein FL *et al.* (1997) Incidence of fetal alcohol syndrome and prevalence of alcohol-related neurodevelopmental disorder. *Teratology* **56**:317–26.
- Shaywitz SE, Cohen DJ, Shaywitz BA. (1980) Behavior and learning difficulties in children of normal intelligence born to alcoholic mothers. *J Pediatr* **96**:978–82.
- Simpson VJ, Rikke BA, Costello JM *et al.* (1998) Identification of a genetic region in mice that specifies sensitivity to propofol. *Anesthesiology* **88**:379–89.
- Smolen A, Marks MJ, Smolen TN *et al.* (1986) Dose and route of administration alter the relative elimination of ethanol by long-sleep and short-sleep mice. *Alcohol Clin Exp Res* **10**:198–204.
- Sokol RJ, Delaney-Black V, Nordstrom B. (2003) Fetal alcohol spectrum disorder. *JAMA* **290**:2996–9.
- Sorensen S, Palmer M, Dunwiddie T *et al.* (1980) Electrophysiological correlates of ethanol-induced sedation in differentially sensitive lines of mice. *Science* **210**:1143–4.
- Steinhausen HC, Spohr HL. (1998) Long-term outcome of children with fetal alcohol syndrome: psychopathology, behavior and intelligence. *Alcohol Clin Exp Res* **22**:334–8.
- Steinhausen HC, Willms J, Spohr HL. (1993) Long-term psychopathological and cognitive outcome of children with Fetal Alcohol Syndrome. *J Am Acad Child Adolesc Psychiatry* **32**:990–4.
- Stoler JM, Ryan LM, Holmes LB. (2002) Alcohol dehydrogenase 2 genotypes, maternal alcohol use, and infant outcome. *J Pediatr* **141**:780–5.
- Streissguth AP, Dehaene P. (1993) Fetal alcohol syndrome in twins of alcoholic mothers: concordance of diagnosis and IQ. *Am J Med Genet* **47**:857–61.
- Streissguth AP, Herman CS, Smith DW. (1978) Intelligence, behavior and dysmorphogenesis in the fetal alcohol syndrome: a report on 20 patients. *J Pediatrics* **92**:363–7.
- Thomas JD, Goodlett CR, West JR. (1998a) Alcohol-induced Purkinje cell loss depends on developmental timing of alcohol exposure and correlates with motor performance. *Dev Brain Res* **105**:159–66.
- Thomas JD, Leany BD, Riley EP. (2003) Differential vulnerability to motor deficits in second replicate HAS and LAS rats following neonatal alcohol exposure. *Pharmacol Biochem Behav* **75**:17–24.
- Thomas JD, Melcer T, Weinert S *et al.* (1998b) Neonatal alcohol exposure produces hyperactivity in high-alcohol-sensitive but not in low-alcohol-sensitive rats. *Alcohol* **16**:237–42.
- Ulug S, Riley EP. (1983) The effect of methylphenidate on overactivity in rats prenatally exposed to alcohol. *Neurobehav Toxicol Teratol* **5**:35–9.
- Viljoen DL, Carr LG, Foroud TM *et al.* (2001) Alcohol dehydrogenase-2*2 allele is associated with decreased prevalence of fetal alcohol syndrome in the mixed-ancestry population of the Western Cape Province, South Africa. *Alcohol Clin Exp Res* **25**:1719–22.
- Vorhees CV. (1989) A fostering/crossfostering analysis of the effects of prenatal ethanol exposure in a liquid diet on offspring development and behavior in rats. *Neurotoxicol Teratol* **11**:115–20.
- Vorhees CV, Fernandez K. (1986) Effects of short-term prenatal alcohol exposure on maze, activity, and olfactory orientation performance in rats. *Neurobehav Toxicol Teratol* **8**:23–8.
- Wainwright PE. (1998) Issues of design and analysis relating to the use of multiparous species in developmental nutritional studies. *J Nutr* **128**:661–3.
- Webster WS, Walsh DA, Lipson AH *et al.* (1980) Teratogenesis after acute alcohol exposure in inbred and outbred mice. *Neurobehav Toxicol* **2**:227–34.
- Westergren S, Rydenhag B, Bassen M *et al.* (1996) Effects of prenatal alcohol exposure on activity and learning in Sprague-Dawley rats. *Pharmacol Biochem Behav* **55**:515–20.
- Wigal T, Amsel A. (1990) Behavioral and neuroanatomical effects of prenatal, postnatal, or combined exposure to ethanol in weanling rats. *Behav Neurosci* **104**:116–26.
- Wozniak DF, Hartman SE, Boyle MP *et al.* (2004) Apoptotic neurodegeneration induced by ethanol in neonatal mice is associated with profound learning/memory deficits in juveniles followed by progressive functional recovery in adults. *Neurobiol Dis* **17**: 403–14.
- Yanai J. (1983) Genetic factors in drug neuroteratogenicity. *Subst Alcohol Actions Misuse* **4**:19–30.
- Yanai J, Ginsburg BE. (1977) Long term reduction of male agonistic behavior in mice following early exposure to ethanol. *Psychopharmacology* **52**:31–4.
- Yanai J, Ginsburg BE. (1979) The relative contribution of pre- and neonatal ethanol administration to changes in mice behavior. *Arch Int Pharmacodyn Ther* **241**:235–44.
- Zimatkin SM, Deitrich RA. (1995) Aldehyde dehydrogenase activities in the brains of rats and mice genetically selected for different sensitivity to alcohol. *Alcohol Clin Exp Res* **19**:1300–6.
- Zorilla EP. (1997) Multiparous species present problems (and possibilities) to developmentalists. *Dev Psychobiol* **30**:141–50.